

: 1

28	1.1
28	1.1.1
31	2.1.1
34	3.1.1
36	2.1
44	3.1
49	4.1
56	5.1
59	6.1

WHO

. "Model Prescribing Information. Drugs used in Anaesthesia. Geneva: WHO; 1989

(intubation) ()

2.10 ()

) 7.18 () 1.3.18 () 1.18 ()

.(

1.1

1.1.1

7-4

15

Thiopental sodium

1 0.5 ()

:

:

:

.1 : (2) (5)

24

24

:

150-100 15-10 (25) %2.5

60 150-100 ()

60 7-2 4

20 25 .

24 . 1 40 0.5

:

Ketamine

10 1 50 () ()

:

() :

) :

(

1 : (2)

24

:

10)

13-6.5

(25-12

2)

4.5-1

(10-5

1

45-10 ()

2 0.5

4

)

:

(24

2.1.1

()

Ether, anaesthetic

(1988)

() :

:

1 : (2) :

:

%15

%10 () %5-3

:

Halothane

:

:

)

:

18

(

1 : (3 2)

()

:

%2-1.5 () % 4-2

()


%2-0.5

:

3.1.1

) (5.12) (1.3)

(1.25



Nitrous oxide

() :

:

.1 : (2) :

:

%30-25

%50

%50

:

Oxygen

:

.

()

1 : :

:

%21

%80

:

2.1

()

4

%0.5

7

()

)

(3.21

%1

%0.5

%0.5

%5

Bupivacaine hydrochloride

5

10

(%0.25)

2.5

()

75 (%0.5) 5 10 (%0.5)

4 (%7.5)

:

:

(5)

:

1 : (3 2)

:

(60) 150 %0.25

(30) 150 %0.5

(3.6-1.8) 18-9 %0.5

(20-10) 100-50 %0.5

) 60 () %0.5-0.25

(12

30) 150 %0.5-0.25

(

20) 100 () %0.5-0.25

(

1.5 %0.25

:

Lidocaine hydrochloride

10	20	(%0.5)	5	()
75	2	(%5)	50	20 (%1)
				(%7.5)
5	(%1)	10	()	
		20	(200000 1)	
(%2)	20	()	()	

2.2 (80000 1) 12.5

(%4-2) 40-20

:

(2.12)

:

() (5) :

(2) () ()

1 : (3)

:

50) 250 %0.5

(

(25) 250 %1

(5-1) 200-40 %4

(10) 400 %4

(5.1-1) 75-50 (%7.5) %5

) 400 %0.5

(80

) 400 %1

(4

(5-1) 100-20 %2

4 %1 %0.5 :

7 (200000 1) 5 + %1 %0.5

:

()

)

.(

Ephedrine hydrochloride

1

30

()

:

(4)

:

1 : (3 2)

:

3

4-3

(9)

6-3 ()

30

()

:

Epinephrine (adrenaline) ()

:

:

:

1 :

:

(20000 1)

5

500

(80000 1)

12.5

3.1

(5.1)

()

Atropine sulfate

1

600

()

:

(5.17)

(3.2.4)

(5.21)

:

:

:

1 : (3 2)

:

:

20

60-30

500

30-10

1-0.4

20

1.2-0.6

3-2

:

()

Diazepam

(1971)

5

2

2 5 ()

:

()

(3.24) (1.5)

:

:

(4) (5) ()

1 : (3 2)

24

.

:

10-5 12

200 12

:

Promethazine hydrochloride

	25	10	
	5	5	()
2		25	()
	(2.17)		:
			:
2)	(5)		:
		1	: (3
24		.	
			:
	1-0.5		
	25		
		()	:

()

()

()

30

(30-20)

.2.20

2 5 ()

:

:

(5 4) :

1 : (3 2)

:

50 250-200

50 200-125

:

()

Vecuronium bromide



10 ()

:

:

(4) :

)

1 : (3 2) (

:

100-80 5

20-10 4

30-20

1.4-0.8

100-40

) :

Suxamethonium chloride

2 50 ()

()

:

:

:

()

()

:

)

()

: (2) (5) (

1

:

4

5-4

150

1

2

10-5

1-0.5

4-2.5

()

500

2-1

:

)

(

(

)

()

Neostigmine metilsulfate

2.5

1

500

()

1

:

(2. 20)

:

)

:

(

(4)

(

)

1 : (3 2)

:

2.5

40

5

500

()

1.2-0.6)

:

(20

) 500

(

:

5.1

()

2.2

.1.2

Morpine

(1961)

1 10 () ()

) :

(2.2

) :

(5

:

(2) (4)

1 : 2.2.4 : (3)

:

200-150

100-50

60-40

100

-100

4

300-150

2.5-2

30

10-8

200

:

Naloxone hydrochloride

1

400

()

)

:

(2.2.4

:

:

3-2

200-100

100

10

10

:

%15

70

9)

(%0.9

(%0.18 %4)

24

.() 26.2 () 11.1